

Common Rapid Response Team Interventions Post-Test

- 1) In what decade did the idea of rapid response teams develop?
 - a) The 1960's
 - b) The 1970's
 - c) The 1980's
 - d) The 1990's

- 2) An oxygen saturation less than _____, despite supplementation, is a valid reason to activate a rapid response.
 - a) 88
 - b) 89
 - c) 90
 - d) 91

- 3) A patient experiencing seizure-like activity, while positive for benzodiazepines, should be given Flumazenil as an antidote.
 - a) Yes
 - b) Yes, but they should also be given Narcan
 - c) No, they should be given Narcan instead
 - d) No

- 4) For severe alcohol withdrawal the best course of action is:
 - a) Benzodiazepines or antipsychotics pushes
 - b) Benzodiazepines or antipsychotics IV drip
 - c) Aerosolized alcohol
 - d) No treatment, just wait it out

- 5) Blood sugar levels less than 40 mg/dL are considered:
 - a) Normal
 - b) Mild hypoglycemia
 - c) Severe hypoglycemia
 - d) Mild hyperglycemia

Common Rapid Response Team Interventions Post-Test

- 6) Hyperkalemia is classified as:
- a) Potassium level greater than 5.5
 - b) Potassium level less than 4
 - c) Chloride level greater than 115
 - d) Chloride level greater than 140
- 7) Hyperkalemia might cause which ECG changes:
- a) Flattened T waves
 - b) Widened P waves
 - c) Narrow QRS
 - d) Peaked T waves
- 8) A trach that is 4 days post placement has come out. What is the preferred course of action?
- a) Attempt to put it back in
 - b) Wait and see if the patient really needed it anyway
 - c) Orally intubate and send to O.R. for re-insertion
 - d) Intubate through the stoma with an ET tube to keep the stoma open
- 9) For a patient with a suspected ischemic stroke you should get a CT or MRI done and read within how many minutes?
- a) 15
 - b) 25
 - c) 45
 - d) 60

Common Rapid Response Team Interventions Post-Test

10) For a patient with an asymptomatic < 15% pneumothorax the best course of action is?

- a) 100% oxygen and observe
- b) Needle decompression
- c) Chest tube
- d) 100% oxygen and CPT